
### A healthy outside, Starts from the inside.



### A healthy outside, Starts from the inside.



144 800mg

98 750mg

18 650 mg

131 600 mg

40950 mg

175 mg / kg

10 000 mg

1 315 mg

Composition Each 1 kg

Coriander extract (CoE 154)

Extract from the cinquefoil

Bentonite (lm558)

BHT (E321)

Diallyl disulfide

Extract from oak bark (CoE390)

Polyoxyethylene castor oil (E484)

Carboxymethylcellulose (E466)

### Indications:

- → Binds substances in the gastrointestinal tract, such as bacterial toxins, medicines, and intestinal gasses
- ✔ Increases intestinal peristalsis, thus supports intestines function

### Benefits:

### Coriander

- Strong detoxifying properties.
- Antibacterial effect.
- · A strong antioxidant and a valuable source of vitamins.
- Binds toxic heavy metals and removes all toxins from the body.

### Oak bark:

- Thanks to its astringent properties, it helps in the treatment of diarrhoea (tannins combine with proteins of the intestinal mucosa and cause shrinkage).
- · Has an anti-inflammatory, anti-haemorrhagic, antibacterial and fungicidal effect.

### Potentilla:

- Inhibits compounds resulting from the decomposition of dead cells and damaged tissues, e.g. heavy metals, toxins.
- Prevents minor bleeding from damaged capillaries in the mucous membrane of the entire gastrointestinal tract.
- · Reduces inflammation of gastric and intestinal mucosa.
- · Prevents dilation of blood vessels.

### Activated carbon:

- Strong adsorbent (binding) properties towards other substances, including bacteria and toxins causing diarrhoea. Captures and binds them and is then excreted from the body with them.
- Covers the mucous membrane of the digestive tract with a layer protecting it from harmful substances.

### Dosage:

Cow Buffalo: 100 g of preparation 2 times a day.

Calves: 20 g of preparation 2 times a day.

Lambs: 2-5 g of preparation 2 times a day.

### **Storage Conditions:**

- · Store in a cool and dry place at room temperature.
- · Keep out of reach of children.
- · Use as per veterinarian's recommendation.



or Veterinary use only

Manufacturer:

• بدآنتوں کی حرکت کودرست کرتا ہے۔ • بدنظام باضمه کودرست ہے تا کہ مادوں کا اخراج ہوسکے۔ گے بھینس:100 گرام دن میں دوبار مر عن 20 گرام دن میں دوبار مره، ديوفيره: 2 سے 5 گرام دن ميں دوبار

. برمعدے میں موجود بیکٹر بل زہر یلے مواد، ادویات،

فاضل مصنوعات، زہریلی گیس وغیرہ کوصاف کرتاہے۔


• كرے كے درجة رارت يرخشك جله يروكيس-. بچول کی پہنچ سے دورر کھیں۔

• وٹرزین کی ہدایت کےمطابق استعال کریں۔

Exclusive Distributor:



# Refuel the energy that moves Lives





### Avoiding Problems before they happen Mitigate the negative energy balance effect

### **Ketosis! What?**


A common condition - affecting around 30% of cows


### Ketosis..... So What?

Consequeces: Imapcts cow health, fertility
and milk production
After effect: Expensive, stressful and cow welfare

### **Ketosis..... Now What?**

**ENERGY**- a novel solution to reduce the incidence of ketosis





Negative Energy Balance: An imbalance between dry matter intake and production needs.



### **Avoiding Problems before they happen** Mitigate the negative energy balance effect

Compromised energy state compounds result in mobilization of fat in the body. Ketone bodies increase in the blood, resulting in subclinical ketosis. Subclinical: 1.2-2.9 mmol/L Clinical:> 3.0 mmol/L

Increased immunosuppression+ decreased DMI makes the herd more susceptible for secondary diseases and infertility issues.

Short term: Metritis, mastitis, removal from herd, DA Long term: less likely to reproduce within 120 day cycle, 700 lbs. less lactation, fatality

### Secondary Disease



#### Metabolic Demand

Doubles as early as 2 days post-calving and sends the cow into a metabolic compromised state.

### **Immunosuppression**

With increased ketone bodies, inflammatory levels in the body increase and results in immunosuppression. This reaction also decreases DMI.

Glycerol, with its high and rapidly available energy, has been used for many years as a supplement to alleviate ketosis in dairy animals (Cotrill et al., 2007, Donkin, 2008).

With low feed intake and glucose supply, the dairy cow mobilizes large amounts of body fat, causing fat accumulation in the liver that results in lowered blood glucose.



### Very High Energy

Glycerine and Dextrose give a rapid energy boost to cows when they need it the most.



### **Combat Negative Energy Balance**

Avoids Negative Energy Balance to reduce post calving metabolic diseases such as Ketosis and improve fertility rates.



### **Improved Intakes**

Energy Drink will help boost post-calving intake levels, helping milk production as well as liver function



### Easy To Use

Ideal Use via dispenser





### Avoiding Problems before they happen Mitigate the negative energy balance effect

- Reduces the incidence of Ketosis.
- Easy to administer.
- Supports regeneration after calving.

### **Components:**

Glycerine, Dextrose

### **Additives** Per 1Liter

Vitamin B3	10 000mg
Potassium iodide	500mg
Vitamin B2	400 mg
Vitamin B9	166mg
Vitamin B12	. 12,6mg
Cobalt carbonate	. 8mg

### **Preservatives:**


• Calcium propionate 2000mg

### **Sensory Additives:**

• Mixture of flavoring compounds 100mg

### How to use:

Administer orally immediately after birth via dispenser. Daily dose is 500ml. Recommended use for 3 days after calving.










### FARMIX-AR Multidimensional Feed Presentation of Multidimension o

### Multidimensional Feed Premix / Feed Grade

For High levels of milk production, Energy requirement are met by encouraging maximum intakes of energy dense, low fiber diet with highly fermentable carbohydrates.

This re



### **FARMIX-AR**

Multidimensional Feed Premix / Feed Grade Vitamins, Minerals & Yeast

Farmix-AR is a premix feed that must be added into a feed of animals in order to achieve the sufficient level of live yeast, vitamins and minerals that prevent the animals from malnutrition, deficiencies, and acidosis, maintain rumen ph. and increase milk production.

فا کس اے۔ آر اپنی اجزائے ترکیبی میں موجود وٹا منزی بدولت جانور کیلئے متوازن خوراک کی بنیا و فراہم کرتا ہے۔ فارکس اے۔ آر اجزائے ترکیبی میں موجود منزلز کی بدولت، میٹا پولک افعال کے فروغ سے پیداوار کی صلاحیت میں برحوتی اور توت میں اضافہ اور تولیدی صلاحیت میں اضافہ بوتا ہے۔ فارکس اے۔ آرمیں وٹا من Eliological cell membrane کی صوجود گی حیاتیاتی غلیہ (Intracellular antioxidant) کی سمائیت کی ضامن ہوتا ہے۔ فارکس اے۔ آرمیں دو Yeast (خمیر ) کا بڑوکا معدہ (Rumen) کے کار آمداور مفید جرثو مول کوخوراک فراہم کر کے معدہ (Rumen) کی میٹا پولام میں اضافہ کرتا ہے۔ جانور کی اجزا کا معدہ کی مسامنہ کی سمائی کی بیٹا پولام میں اضافہ کرتا ہے۔ جانور کی جداوار کی سلاحیت اور دودھ کی پیداوار میں اضافہ کرتا ہے۔ نیٹ میں کی واقع ہوتی ہے۔

### : Biotin

Biotin کی موجود گی گھر ول کی صحت اور دودھ کی بڑھاتی میں اہم کر دارا داکرتی ہے۔

### : Niacin

فارکس اے آریس Niacin کی موجود گی Ketosis کو کم کرنے اور نظام ہضم میں acidosis کو کم کرنے اور نظام ہضم میں در ادر جانورکو rotozoan) Entodinia ( تیزابیت ) سے محفوظ رکھتی ہے۔

### منرلز (Minerals):

فارکس اے آرمیں مائنگر ومنرلز

اور (Zinc, Selenium, Manganese, Iodine, Copper, Cobalt) اور (Calcium, Magnesium, Sodium, Phosphorus) کی موجود گی تو سدا فعت ، دو ده کی پیدا دار ، جسمانی بره حوتی ، تو لیدی صلاحیت بیس اضافه کا سبب ہیں۔ مثلاً Copper اور Copper کی موجود گی تو لیدی صلاحیت میں اضافه کرتی مشافه کرتی د retained placenta اور early embryonic death کے نیز Pitamin B اور Cobalt کے فار مزکوم حاثی فقصان سے بیجاتی ہے۔ Cobalt جسم میں Osmotic pressure برقو ہے۔ منر لڑجسمانی Osmotic pressure برقو ہے۔ منر لڑجسمانی صدر گار ہے جو بہت اہم بڑو ہے۔ منر لڑجسمانی Cosmotic pressure برگھتے ہیں۔

کیلیم اور فاسفورس ہڈیوں کی نشونماء میں اہم ہیں اور ہڈیوں میں اِن کی موجود گی جانور کو کسی بھی فوری کی سے محفوظ رکھتی ہے۔ میکنشیم کا کاریو ہائیڈریٹ کی میٹا پولزم میں اہم کردار ہے۔ سوڈیم اور کلورین pH ، Cellular Volume

### فارکس اے - آر کے اجز ائے ترکیبی کے فوائد: خمیر (YEAST) :

نظام بمضم کی pH کومتوازن رکھ کرموافتی ماحول فراہم کرتا ہے تاکہ باضمہ (Rumen) اپنے افعال بہتر مین طریقہ سے انجام دیاور جسمانی بردھوتی اور دود دھ کی مقدار میں خاطر خواہ اضافہ ہو۔ خوراک میں موجود Proteins, Fats کو ہیں کہ اور Carbohydrate کو ہاضمہ Fiber کو جیس کرسکتا۔ فار کس اے۔ آرکے دوخمیر (Yeast) کو جنب کرسکتا۔ فار کس اے۔ آرکے دوخمیر (digest کو Fiber کو Lactic Acid کرنے میں مددگار ہیں جس سے دود دھ میں اضافہ ہوتا ہے۔ لیکھوٹا مول کی تعداد بردھا کر Lactic Acid کی پیداوار کو کم کرتا ہے۔

Rumen پر Stress اورگرمی سے پیداشدہ اثرات کو کم کرتا ہے اور پڑھوتی اور وودھ کی کی پیداوار کو متارث نہیں ہونے دیتا۔

### وٹامن (Vitamin):

فارکس اے- آرمیں موجود وٹامن جانوروں میں گئی بیار یوں کا ندارک کرتے ہیں۔وٹامن متوازن خ<mark>وراک کی اہم ترین بنیاد ہیں۔</mark>

وٹامن اے (Vitamin A): ہڑیوں کی نشو دنما ، قوت ما فعت ، جلداور کھروں کے نشوز کیلئے مفید ہے، نظر کیلئے مفید ہے۔ سانس اور تو لیدی مسائل کے خلاف کار آمد ہے۔

وٹامن Vitamin D3)D3): سوتک کے بخار کے خلاف موثر ہے، ہڈیوں کی بناوٹ کیلئے ضروری ہے۔ جانور میں کیلئیم فاسفورس بیلنس کو برقر اررکھتا ہے۔

وٹا من ای (Mastitis :(Vitamin E) (ساڑو) کے خلاف قوت مدافعت بناتی ہے۔ وٹا من E اور سیکنیم کی موجودگی tissue degeneration (خلیوں کے انحطاط) مثلاً کی روک تھام کے خلاف مفید ہے۔ گوشت میں لال رنگ (Oxymyoglobin) کو برقر اررکھتی ہے۔

وٹامن لیEnergy Metabolism(Vitamin B12)12 اور

### **خوراک:** 0.5 سے 1 کلو 1 ٹن فیڈ کے لئے گائے: 30-50 گرام روزانہ 100-50 گرام ( بیاری کی صورت میں ) بھیڑاور ب**کری:** 10-20 گرام روزانہ

قوت مدا فعت برقرارر کھنے میں مددگار ہے۔

بچھڑااور میمنے: 5گرام روزانہ

#### **Exclusive Distributors:**



### Dosage:

0.5 kg to 1 kg per ton of feed.

Cow: 30 - 50 gm per day (In routine)

50 - 100 gm per day (Stress / diseased conditions)

**Sheep & Goats: 10-20** gm per day. **Kids and Lambs: 5** gm per day.


### Manufactured by:



Balance



# Balance



### Composition Each 1 Liter


190mg Calcium D-pantothenate (3a841) Vitamin B1 HCI (3a820) 180mg Vitamin E (3a700) Vitamin B2 400mg 5 000 IU Cobalt sulphate 7H20 (3b305) Vitamin B3 (3a315) 15 000 mg 10mg Vitamin D3 (E671) 4 000 000 IU Vitamin B6 (3a831) 170mg Orthophosphoric acid 85% (1a338) 1,625 mg/l Vitamin B12 60mg

Feed Materials: Glycerine (13.8.1), Sucrose (4.1.3), Calcium Chloride (11.1.6), Magnesium chloride (11.2.6)





### Different mineral deficiencies with downer cows




- Ca-deficiency
- Combined Ca-and P-deficiency
- P-deficiency
- Ca and P in normal state
- Mg- deficiency

Prophylactic oral administration of Phosphorus and Calcium can reduce the development of hypophosphatemia and milk fever at calving

<sup>6</sup> Pichon, S: Analyse von Festlegenursachen be i kuhen - Eine Praxisstudie. Disseration, Leipzig, 2008

Milk fever and subclinical hypocalcaemia are the most important macro- mineral metabolic disorders that effect transition dairy cows.



Sub-clinical Cases

Clinical Cases

For every case of clinical milk fever (visible) you have, you are likely to have 4 → 6 cases of sub-clinical (invisible) cases

Incidence of clinical hypocalcaemia is approximately 5%. However, incidence of sub-clinical hypocalcaemia (SCH) has increased upto 78%

78%
CAN HAVE
SCH

(Reinhard et al., 2011)





Most of the cases of milk fever occur within one day of calving because milk and colostrum production drain calcium from the blood





Occur within 24 hours after calving

3% Cases

Occur few days before calving

6% Cases


Occur just few hours before or at

time of parturition


4% Cases

Occur 48 hours after calving

Diseases related to immune suppression and energy balance in the transition dairy cows




Milk fever will substantially increase the risk of many other problems by manyfolds.




# Balance

8924 - 3 347

Subclinical hypocalcaemia increased with age and was present in 41%, 49%, 51%, 54%, and 42% of 2nd – 6th lactation of cows, respectively




Calcium is critical for proper immune cell function. Hypocalcaemia decreased a number of neutrophils which increased the risk of infectious diseases.





Cows with low plasma vitamin E levels before calving have a 9 times higher probability of clinical mastitis as compared to cows with an acceptable blood level.

NRC in 2001 recommends higher dietary vitamin E levels for transition and lactating cows to maintain the minimum plasma vitamin E concentration for optimum immune response.



Calcium supplemented orally (not part of the diet) after calving has shown a positive response for preventing a drop in concentration of blood calcium.






### **Immunity Booster for Calves After Birth**

- Highly concentrated dose of antibodies




## Rear Better Calves

Causes of death in the first week of the Calve's Life




The time when a calf is transitioning from passive immunity derived from its own active immunity during its first week of life are critical.

### Causes of ealf Mortality



More than half of all calf deaths in the first few weeks of life are related to scour.

### Percentage of causes of calf scour



Among different Pathogen causing calf diarrhoea, Rota Virus and Corona Virus cause 37% and 28% respectively




### Rear Better Calves


Rotavirus is the most common causes of diarrhoea in neonatal calves between the ages of 4 and 14 days



Calves (days old)



<sup>^</sup> Other infectious organisms, such as Coccidia, BVDV, Clostridium Perfringens and other Ecoil, may also affect calves, often after 3-4 weeks of age.




Loss of bodyfluid due to dehydration in diarrhoea increases the chance of mortality.

Igy antibodies against Bovine Rotavirus can be used to prevent virus associated diarrhea in newborn calves

(Research in Veterinary Science September 2010)

Rota virus and Corona virus destroy the gut lining of the small intestine reducing the digestive and absorptive capacity causing diarrhoea.







# Rear Better Calves



### **Composition:**

Corn Oil, Coconut Oil, Whey Powder Egg Protein, Dried Bovine Colostrum

### Additives per 1 Liter:

- Vitamin E (3a700) 1000 mg
- Vitamin A (3a672c) 260 mg
- Vitamin D3 (E671) 2 mg

### Preservatives:

- Colloidal silica (E551b) 100 000 mg
- BHA (E320) 150 mg

### How to Use:

Calves, foals and camels: 30ml (one syringe) onto the back of

the tongue immediately after birth; ideally within the first 2 hours after birth.

Repeat the dose after 12-14 hours if necessary.



Components	Benefits
Coconut oil	<ul> <li>An energy source which consists mainly of medium chain fatty acids.</li> <li>Medium chain fatty acids are absorbed directly from the small intestine into the liver, where they are converted into energy components.</li> <li>Passive absorption of fatty acids allows the rapid release of energy components for direct use by calves.</li> </ul>
Egg whites	<ul> <li>Help strengthen the passive immune systems.</li> <li>Provide active proteins which counteract, rota viruses and corona viruses.</li> <li>Support natural immunity to common diseases and reduce calf mortality.</li> </ul>
Dried bovine colostrum	<ul> <li>Helps strengthen the passive immune system.</li> <li>Provides active bovine proteins.</li> <li>Supports natural immunity to common diseases.</li> <li>Reduces calf mortality.</li> </ul>
Vitamins A, D and E	<ul> <li>Raise vitamin levels in newborn calves.</li> <li>Strong antioxidants.</li> <li>Help improve the immune system response and immunity to diseases.</li> </ul>

**Exclusive Distributor:** 



جانوروں میں استعال کے لئے For Veterinary use only

