

# **THE STANDARDS AND PROCEDURE FOR BREEDING ANIMALS, 2017**

## **LIVESTOCK BREEDING SERVICE AUTHORITY, SINDH**

### **LIST OF CONTENTS**

<b>Chapter No.</b>	<b>Subject</b>	<b>Page No.</b>
1	Standards and Procedures for Performance Recording and Progeny Testing	2-6
2	Standards and Procedures for Selection of Breeding Males for Artificial Insemination	7-9
3	Standards and Procedures for Selection of Breeding Males for Natural Services	10-13
4	Standards and Procedures for Physical Infrastructure for Semen Production Facilities	14-17
5	Standards and Procedures for Semen Collection and Preservation Standards	18-22
6	Guidelines and Standards for Import of Exotic Semen	23-24
7	Standards and Procedures for Artificial Insemination Training and Training Institutions	25-29
8	Guidelines and Standards for use of Exotic Semen	30-32
9	Standards and Procedures for Artificial Insemination Services	33-34

# **THE STANDARDS AND PROCEDURES FOR BREEDING ANIMALS, 2017**

## **LIVESTOCK BREEDING SERVICE AUTHORITY, SINDH**

### **NOTIFICATION**

In exercise of the powers conferred under section 8, 9,10,11 and 12 of chapter- III of the Sindh Livestock Breeding Act 2017 vested to the Livestock Breeding Service Authority and now the said authority is pleased to make the following standards and procedures of breeding:-

1. Short title, extent and commencement.-These standards may be called the Standards and Procedure for Breeding Animals, 2017 and they shall come into force at once.

### **2. I: Chapter 1**

#### **Standards and Procedures For Performance Recording and Progeny Testing**

Performance recording is basic requirement for any progeny testing program. It includes economically and aesthetically important qualities. Progeny testing is utilized to select males/bulls for genetically controlled traits/qualities of economic importance such as milk production) for which they possess genes but they do not exhibit those traits. Successes in artificial insemination and embryo transfer technologies in dairy cattle and buffaloes necessitate that bulls with better genetic potential be disseminated widely. To judge that a bull has better potential for traits which are not expressed in the males, female relatives are evaluated. When a bull calf is born, information on its mother and other ancestors (e.g., grand mother) and even sisters (if they had performance records) may be utilized for such a judgment. Such a decision making is called pedigree or family selection. The problem with such a decision is that such an information is very limited (e.g., few lactations in few herds) and therefore level of accuracy in declaring a calf good or bad may not be very high. As such a male can have a very profound influence on the population, level of accuracy needs to be much higher to use it extensively.

Information on the performance of large number of daughters can help to really estimate whether a bull has good potential for traits like milk yield and such a selection is therefore called progeny testing. Nowadays, availability of computers have made it possible to utilize all available information on the relatives (pedigree, progeny and other relatives) of a bull and therefore it is not just the progeny that contributes to the evaluation of a bull. Still we may call it progeny testing instead of animal model evaluation or some other kind of evaluation, for simplicity.

## **Breeding Objectives**

Before taking up the progeny testing program, breeding objectives i.e main purpose of keeping that particular breed should be clear. It may be combination of few or many traits. For Red Sindhi cattle, Kundhi buffaloes and Kamori for example, progeny testing is being done for milk yield while traits like fat yield, type/beauty etc are only rarely considered. For other Livestock breeds of Sindh like Tharparkar, Kankrej, Pateri, Tapri etc it may be meat with milk yield as secondary trait. The procedures outlined are therefore focused on progeny testing for milk production.

## **Steps required**

### **1. Survey of the area for identification of potential farmers:**

Identification of potential farmers to be involved in progeny testing program. Bigger farmers (>10 breeding females of the breed) that are willing to cooperate, in terms of actual pedigree and performance recording and use of 100% A.I is needed. Agreement should delineate the services provided to the farmer and information to be collected from him. It should for example, specify that he will not keep any bull and that about half of the females will be inseminated with young bulls and other half with proven bulls.

### **2. Training of AI technicians and recorders**

Milk recorders and artificial insemination technicians recruited for the project need to be trained /made available. Milk recorder should at least Matric science group, with good social skills.

### **3. Tagging and registration of cows/buffaloes**

Animals should be tagged or freeze branded to have a unique identification. A digital photo is nowadays quite cheap. A closer photo of head to capture side face and horns, eyes and face should supplement the tag/brand. About 200 pregnancies per bull are intended (and assuming a calving interval of 1.5 years and two inseminations per conception) it would mean that 600 cows/buffaloes per bull need to be identified for getting preliminary evaluation of a bull. To test 100 bulls therefore 50-60 thousand females are needed to be registered. All the animals of the breed with the registered farmer/herd need to be tagged and recorded. No minimum milk production level is required. As all the animals served with natural service bulls and through Artificial Insemination also need to be tagged. Ear tag should be unique and should have district code and animal number. For example a Kundhi buffalo in Hyderabad district may have KUB00001HYD. This will be the same number as registration number with the Authority. AI technicians will also be using the same number even when animal is not under performance recording program.

### **4. Performance recording**

Milk recording of all the cows/buffaloes in any registered farm is needed as per international norms, not just good/elite females. For milk yield, morning and evening recording at approximately monthly interval is an acceptable frequency. When cows/buffaloes are milked in the presence of calf, milk consumed by the calf (e.g, one teat) may be appropriately accounted for the recorded production or calf may be fed separately and cow/buffalo recorded completely. Lactation yields should be projected using Test Interval Method ([www.icar.org](http://www.icar.org)). Multiple

lactation records should be transformed into an index compensating for repeatability of the trait. Other than milk and fertility records, health status also needs to be recorded. Beauty may be an important trait and prospective bull mothers should be above average in beauty standards, especially the udder and feet-and-legs. Supervision of performance recorders is needed and accurate work should be recorded. Planned mating is mandatory for progeny testing programmes.

### **5. Breeding control of registered cows/buffaloes**

All the cows/buffaloes may be inseminated artificially with semen from young bulls and technical experts should make such a choice to ensure randomness in usage of young bulls to be tested. Record of every insemination, bull used, pregnancy and birth will be kept on a specific recording sheet.

### **6. Processing of recorded information**

The recorded data on farmers cows/buffaloes may be edited monthly after they are received from the field, may be edited for consistency and for generation of reports for feed back to the farmers and for editing to check if it conforms to the already available information of the animal and the farmer. Although such processing of information may be done routinely, generation of reports for feedback may be done quarterly (every three months) and genetic evaluation done biannually if not at a more frequent interval. Lactation yields need to be predicted from monthly records and decision for any cow/buffalo to be elite or not elite needs to be made and communicated to the relevant field staff for planned breeding if needed. Choice of lactation length may be 10 months (both for cattle and buffaloes) to conform to international standard. Lactations ending normally may not be projected to any standard lactation.

### **7. Identification of elite cows/buffaloes**

Elite cows and buffaloes are routinely declared and there is no minimum level of milk production for such a declaration. Generally these are the best animals within a management group. Their number may depend on the number of bulls to be tested every year. For example if 500 male calves are intended to be produced to get 200 candidate bulls, 800 births are expected which would require at least 2000 elite cows/buffaloes and if they are top 20% of the recorded population, recorded population will be ten thousand cows/buffaloes. For better selection intensity, recorded population may be even higher. It may be made clear that under good management, elite may a cow/buffalo producing more than 3500 liters a lactation while under ordinary management, elite may be a cow/buffalo producing more than 2000 liters a lactation (adjusted for age/parity, season of calving and other differences). Farms may thus be categorized as good management farms (concentrate feeding possible along with good health care) and mediocre management farms (concentrate feeding rare and grazing mainly offered along with other management issues).

### **8. Feedback mechanism to the farmers**

Under small and medium sized farming system, farmers generally know how their individual animals perform or at least they can easily rank their animals from best to worse. The central idea of feedback is to inform them how their animals perform compared to the animals of other farmers in the area and even across various production systems. This helps them keep involved in the system because selection and culling decisions can be more judicious. Monthly lactation summary reports ranking farmers for average production of their cows/buffaloes and

periodic rewarding of those who excel may help farmers pay attention to recording and improving their management. Culling decisions may also be helped by the experts.

### **9. Selection and raising of bull calves**

The calves born to the elite cows/buffaloes may be monitored carefully based on the breed characters and physical conditions of the calf.

### **10. Selection of candidate bulls**

The bull calves that mature to a breeding age (at the calf raising centers) and donate semen for preservation and later use in the registered herds are called candidate bulls. Their selection is made from among the calves bought to CRCs on the basis of breed characters, libido and semen quality and through advance technology. Sometimes indices are used to rank them. Well-fed young bulls should donate semen at 2 to 2.5 years of age. Age is slightly more in buffaloes than in indigenous cattle. A performance sheet may be used to record performance of bull calves and bulls.

### **11. Semen collection and storage**

The semen production units generally have adequate expertise to train bulls for semen collection and then actual collection of semen, its evaluation and storage after freezing. At least 2 thousand doses should be used to get elite dams (bull mothers) pregnant for getting progeny records and another 8-10 thousands for future use when daughter information becomes available. This means that a bull needs to spend about 2-3 years at the Semen Production Unit. This time can be reduced tremendously by better management at the SPU. After collection of desired doses of semen such bulls can be denoted to the farmers for natural mating especially for farms where Artificial Insemination facilities are difficult to reach.

### **12. Artificial breeding in the registered herds**

The registered herds may be provided with Artificial insemination facility as and when they need it. Preferred method would be a central calling facility where all the Artificial insemination technicians are hooked and nearest one may reach the farm as and when such a request is received (like a taxi service). Periodic monitoring of the system may also be needed to improve efficiency.

### **13. Genetic evaluation of bulls**

The bulls to be ultimately used in Artificial insemination are tested at many stages but final selection is through genetic evaluation for the milk yield (and other traits) i.e estimated breeding values. These evaluations should be done sooner than on yearly basis.

Software to be utilized for genetic evaluation is also important. Programs such as ASREML may be bought or free software such as DFREML or JAA can also do the job. Minimum criteria for any bull to qualify for getting PTAs published may be reliability which may be 50% in the beginning and can be improved gradually to say, 70% and more. Number of daughters to publish any bull's EBVs should not be less than 10.

### **14. Disposal of semen from below average bulls**

Bulls having negative (below average breeding values for the traits like milk yield (or index having other traits) need to be disposed off even when reliabilities on the estimated breeding values are not very high. The bulls if alive may also be disposed off accordingly. At

least bulls ranked at the bottom (below 25%) should be culled and semen discarded.

#### **15. Contractual mating of elites with elites**

The top most bulls (say top 5%) need to mate with top cows/buffaloes to get 200 inseminations for each bull to get bull calves again. The choice of the elites will be from step 13 where genetic evaluation of cows/buffaloes and the bulls is done simultaneously.

#### **16. Continuity of the progeny testing program**

System should be developed in such a way that if a person or two leave, the system does not collapse. So more than one person be trained at every level. One should have capacity to adjust to new technologies and innovations in the field of biotechnology and computational breeding. It may start from simple and feasible data collection protocols and should lead towards an ideal situation of automation with the passage of time.

#### **17. Practicability and authenticity**

Starting any progeny testing program is difficult and maintaining it is even more difficult as at administrative and political level quick fixes have more attraction than any long term endeavor. Recording less may be acceptable but recording false is not. Simplicity helps at least in the beginning and socioeconomic factors may be very important to any tailor-made effort. Sustainability of the progeny testing program is extremely important and requires dedication.

## **2.II : Chapter 2**

### **Standards and Procedures** **For** **Selection of Breeding Males for Artificial Insemination**

Bull is the most important component of any breeding program as everything revolves around it. Bull used for widespread use especially through Artificial insemination must be better than others not just in its abilities to transfer desired characters; it should be free from genetic and other diseases. Best option is that it is progeny tested i.e information on daughter's performance for traits like milk yield was available and estimated breed value(s) calculated. But when/until such bulls are not available, at least it should have performance recorded pedigree i.e it should come from a dam which is better than others in a similar production environment. In progeny testing programs such dams are recorded for traits such as milk yield (morning-evening approximately at monthly interval) for a standard duration of 10 months. Following minimums are to be considered for various breeds and bulls without a recorded pedigree should not be used for semen collection and freezing. Dams are expected to be born and perform under agro-climatic production setup of Sindh. Bulls of indigenous breeds are expected to have been registered with relevant registered associations/societies and pedigrees authenticated. Yet, for breeds where a registered association does not exist, Livestock Breeding Services Authority will issue the breed standards and register such bulls directly for semen production businesses (template for registration of bulls is attached as Annex-1). Unique number will be issued to each bull. They may retain their ear tag numbers and branded numbers.

**Minimum criteria for dam's 1<sup>st</sup> and best lactation yields of bulls brought to semen production facilities.**

<b>Breed</b>	<b>Dam's 1<sup>st</sup> lactation milk yield *(litres)</b>	<b>Dam's best lactation milk yield (litres)</b>	<b>Fat %</b>
Red Sindh	2000	2500	4.5
Thari	1500	2000	4.5
Kankrej	1200	1500	4.5
Kundhi	2500	3000	6.5
Holstein (pure breed)	5000	6500	3.5
Jersey (pure breed)	3500	4500	5.0
Holstein cross breed	4000	5000	4.0
Jersey cross breed	3000	4000	4.5
Nili Ravi	2500	3000	6.5
Gir Cow	2000	2600	4.5
Rathi Cow	2000	2400	4.5

\*in a standard lactation of 305-days

#### **1. Physical Examination**

Introducing any new bulls into a semen production facility requires quarantine measures and a thorough physical examination which will be conducted by veterinary officer (VO), Authorized by the authority to ensure that the bulls do not display clinical symptom(s) of any infection or any contagious diseases and are structurally correct. Structural anomalies such as cryptorchidism, penile frenulum and (moderate to extreme forms of) leg defects (e.g., hocked-in)

or (moderate to extreme forms of) testicular shapes (e.g., asymmetrical testis) are unacceptable even when pedigrees are popular. In bucks and rams obvious mouth defects (under or overshot jaws) should also be avoided. Testicular size may vary with the age of the male yet, very narrow scrotal circumference is neither economical nor desirable. Breed associations may also fix certain minimums for weight, height or other attributes for breeding males so that semen collection facilities can adhere to these parameters. Gait defects may be temporary or permanent and breeding soundness examination (Annex-II) should help avoid undesirable males.

## **2. Genetic abnormalities**

Bulls will be required to have normal karyotype certificate to rule out chromosomal defects. Specific tests may also be conducted for genetically transmitted diseases like Bovine Leukocyte Adhesion Deficiency (BLAD), Citrullinemia and Deficiency of Uridine Monophosphate Synthase (DUMPS) for exotic breeds and their crossbreds. Beta casein variants (A1 and A2) are also becoming important and in future exotic bulls (Holstein and Jersey) and their crossbreds might be screened to only allow A2 bulls for semen production. Our local cattle and buffaloes naturally possess A2 variant, the desired genotype.

## **3. Quarantine tests**

A minimum quarantine period of 60 days is compulsory before bringing new bulls in a semen station (in case of bulls are brought from known sources, 30 days quarantine is sufficient). This period shall be divided in two periods of 30 days during which a series of compulsory examinations and tests are carried out. Only after favorable results from the health control point, the bull shall be admitted to the semen station.

- a) In the quarantine station new bull/s shall be housed for minimum of 60 days in a place which is effectively separated from the facilities occupied by resident bulls and all equipment used in handling, feeding, watering and cleaning the new bulls shall not be shared with the resident herd(s). Quarantine station and the semen production facility can not share the same boundary wall. Both facilities should not share labor as well.
- b) Each new bull in quarantine station will be tested against major contagious diseases before its entry to resident herd e.g. TB, Brucellosis, Campylobacteriosis and Trichomoniasis. All tests shall be done by an accredited institute or disease diagnostic laboratory.
- c) If any bull is found positive for TB, or Brucellosis, it shall be removed immediately.
- d) During quarantine period, the bulls shall be vaccinated against FMD, HS, BQ, Theileriosis and Anthrax. However, vaccinations against bacterial diseases shall be done only if there is an outbreak or prevalence of a particular disease. Level of comfort (shade, feeding, watering etc) will be similar to the donating bulls. Once the quarantine period is over, all bulls shall be introduced to the resident herd. A written consent / certification of designated official / veterinarian will be required before semen collection can be initiated.

## **4. Testing of Bulls**

The resident bull herd would also be required to go through periodical testing and



vaccinations. Testing protocols for bulls against Brucellosis, Campylobacteriosis, Trichomoniasis, Leptospirosis, Tuberculosis, and IBR are given in Annex III to VII. For international trade OIE guidelines ([http://www.oie.int/fileadmin/Home/eng/Health\\_standards/tahm/A\\_index.htm](http://www.oie.int/fileadmin/Home/eng/Health_standards/tahm/A_index.htm)) will need to follow.

## 5. Vaccination Schedule

The bulls shall be vaccinated against FMD, HS, BQ and Anthrax. However, vaccinations against bacterial diseases shall be done only if there is an outbreak or prevalence of a particular disease. To reduce lay off time, the bulls shall be vaccinated on the rest day or the day after completing semen collection.

The semen station in collaboration with local governments shall arrange for carrying out ring vaccinations for all animals against FMD, HS and BQ within a radius of 5 km around the semen station. Vaccinations against HS and BQ shall be carried out in the areas having incidence of these diseases.

### Culling of Bulls and Semen Doses due to Specific Diseases

Disease	Bulls	Fate of Semen doses
Foot and Mouth Disease (FMD)	Retain	Last one month doses to be discarded, refer Annex-VIII
Brucellosis	Castrate & remove	To be discarded since the last negative test
Tuberculosis (TB)	Remove	To be discarded since the last negative test
Campylobacteriosis	Treat and retain	To be discarded since the last negative test
Trichomoniasis	Treat and retain	To be discarded since the last negative test

The semen station must remove bulls (within 48 hours) which are positive for above diseases, poor libido, poor semen quality, incurable lameness, etc. Besides, the semen station shall also cull those bulls which have completed six years of productive period or 2 lac semen doses, whichever is achieved earlier. These conditions will be relaxed for progeny tested (positive) bulls.

## 6. Parentage Confirmation

Parentage confirmation through molecular evaluation for all the candidates bull entering the semen production unit would be compulsory.

## **2.III : Chapter 3**

### **Standards and Procedures** **For** **Selection of Breeding Males for Natural Service**

Males used for providing natural service should have all the basic norms laid down for those used for collection of semen for Artificial Insemination except that their progeny information may not be available. Individual's own performance (for traits like growth) and pedigree performance therefore could be the main information sources. This is especially true for rams and bucks where males have recorded phenotype for traits like growth rate and breeding objectives include growth as selection criteria. Males used for breeding commercially will be registered under the authority and following guidelines are expected to be helpful to provide a level playing field as well as in the improvement of genetic potential of our breeds.

1. Registration with breed association/society.
2. Pedigree Performance Record.
3. Basic phenotypic qualities.
4. Breeding soundness examination.
5. Genetic and venereal diseases.
6. Maintenance of breeding records.

#### **Registration with breed association /Society**

Registration with the Livestock Authority is through the breed associations for those breeds and species for which such farmers/breeders groups exist. For other breeds and species such registration will be by a cell in the Authority headquarters and minimum criteria will be published/declared by the authority.

#### **Basic phenotypic attributes**

For the main local breeds such as Kundhi and Red Sindhi or even for Kamori or Pateri for which semen is available. Associations should come up with minimum criteria for beauty traits apart from the authentication of pedigrees in the form of pedigree certificates.

Kundhi bulls are required to have black color and curled horns (level of curl may vary but tighter is better) along with (white eye balls, partial or complete) characteristics of Kundhi are required Length of body and height are preferred physical attributes. Size of testicles may vary and both single and double pouched scrotums are acceptable yet, symmetry is crucial. Similarly, being hocked-in (rear leg rear view) is not a preferred attribute. Strength of rear legs is actually quite important breeding males of all species as they have to bear all the weight during mounting. Hooves should also be compact.

Red Sindhi bulls are required to have reddish dun color (a variant of red color). Variation of brown shade may vary from greenish red to darker red yet very dark chocolate or very light brown shades are not preferred. Hump and dewlap are characteristically large. Folds are preferred in the dewlap while hump must not be erect and not broken. Longer tails are preferred with black switch. While body spots are not acceptable yet, small (less visible) spots in the

underline may be acceptable. Symmetry of testicles is mandatory and hocked-in bulls are less acceptable. Hooves should not be worn out especially at a younger age.

The Thari breed bulls are whitish gray in color and gray may deepen, particularly on the fore and hind quarters. All along the backbone there is a light gray stripe. The head is of medium size, the forehead broad and flat or slightly convex above eyes: the front of the horns and face are practically on one plane. The skin between the eyes is often wrinkled, the wrinkles running perpendicularly. The eyes are full and bright. The eyelashes are black and there is a small ring of black on the eyelids. The ears are somewhat long, broad and semi-pendulous and face forwards. Horns are set well apart curving gradually upwards and outwards in the same line as that of the poll with blunt points inclined inwards. A small portion of the skin with hairs extends over the base of the horns. The hump in males is moderately well developed, firm and placed in front of the withers. The dewlap is of medium size and the skin is fine and mellow. The sheath in the males is of moderate length, and is semi-pendulous. Shoulders are light and legs are comparatively short, but in good proportion to the body. The hooves are hard and black, of moderate size and have no tendency to turn out.

The Kankrej breed bulls vary in color from silver to gray to iron gray or steel black. Fore quarters, hump and hindquarters are darker than the barrel. The switch of the tail is black in color. The forehead is broad and slightly dished in the center. The face is short, and the nose looks slightly upturned. The strong lyre-shaped horns are covered with skin to a higher point than in other breeds. The ears are very characteristic, being large, pendulous and open. The legs are particularly shapely and well-balanced and the feet small, round and durable. The hump is well-developed and not so firm. The dewlap is thin but pendulous and has pendulous sheaths. Pigmentation of the skin is dark and the skin is slightly loose and of medium thickness.

Gir bulls are distinctive in appearance and a heavy breed. Their body is well proportioned. Their body color is shining red to spotted white although, their body color can vary from pure red to speckles, yellowish red to white with large red spots. Their skin is soft, thin and glossy. They have a big head with prominent bulging shield and a long face. The convex forehead of Gir is the most unusual feature of this breed. Their convex forehead acts as a cooling radiator to the brain and pituitary gland. Their ears are very long and are an excellent fly and insect swatter. Their eyes are hooded and black pigmented and there is lot of loose skin around their eye area. They have very hard feet which is black in color. The horns of Gir cattle are set well back on their heads and thick at the base. Their horns grow downwards and backwards with an upward curve. They have large hump. The average body weight of Gir bull is about 545 kg.

The Rathu bulls are medium sized animal which are usually brown in color with white patches all over the body. Although they are also found in completely brown or black coat with white patches. Their lower body parts are usually lighter in color as compared to the rest of their body. They have medium sized horns, and their horns are curving outward, upward and inward. Their face is broad between eyes and slightly dished. They have long tail, and the dewlap is fine and loose.

In all the breeds, males should have known and performance recorded pedigrees. Performance of dams is crucial and therefore instead of opinion, physical records are preferred. For goats, seeing the dam (if alive) can further help decided selecting of bucks from dams with

good body length and uniform large udders (not large teats).

Same may be true for Red Sindh and Kundhi where udder is the single most important trait to look at. Yet, when performance recorded dams are available, following minimum performance (similar to those of AI males/bulls) of dams is desired. These naturally servicing males will have the same pattern of registration as with Artificial insemination males.

**Minimum criteria for dam's 1<sup>st</sup> and best lactation yields of bulls brought to semen production facilities.**

Breed	Dam,s 1 <sup>st</sup> lactation milk yield*( liters)	Dam,s best lactation milk yield	Fat %
Red Sindh	2000	2500	4.5
Thari	1500	2000	4.5
Kankrej	1200	1500	4.5
Kundhi	2500	3000	6.5
Holstein (purebreed)	5000	6500	3.5
Jersey (purebreed)	3500	4500	5.0
Holstein crossbred	4000	5000	4.0
Jersey crossbred	3000	4000	4.5
Nili Ravi	2500	3000	6.5
Gir Cow	2000	2600	4.5
Rathi Cow	2000	2400	4.5

\*in a standard lactation of 305-days

**Breeding soundness examination**

The objective of BS exam is to identify infertile and subfertile males and promote fertile bulls/bucks. There are many components to a bull breeding soundness examination that can be simplified as follows.

1. **Health status:** This may involve reviewing the health status of the herd of origin and reviewing the results of previous health records, collecting samples and commissioning more laboratory tests, reviewing vaccination and other treatment history and reviewing previous breeding performance (if available).
2. **Physical condition:** Assessing body size relative to age, body condition, conformation, locomotion and noting signs of any infectious or contagious disease. A more detailed examination may then be required particularly to assess the eyes, the teeth, the legs, feet and off-course, the reproductive apparatus. The prepuce will be examined for any evidence of constriction or discharge. The penis will be palpated to ensure that it is freely moveable within the prepuce and that there is no evidence of any abnormal swellings or growths. The testes will be measured for scrotal circumference and palpated to ensure measuring scrotal uniform size and consistency. They should be freely moveable within the scrotum. Any defect observed (orchitis, excessive fat around testes, hernia, cryptorchidism, hypoplasia, etc.) need to be carefully diagnosed. Trans-scrotal

ultrasonography and testicular fine needle aspiration cytology have also been used under advanced setups to diagnose sub fertile males. An internal examination will also usually be carried out to palpate the accessory reproductive apparatus and ensure normality.

3. **Semen examination:** Electro-ejaculator is commonly used for getting semen sample yet, teasing the bull with a cow that is in oestrus and then using an artificial vagina may be equally good. An assessment can be made to ejaculate volume and density before the sample is viewed using a microscope to assess sperm motility and the number of abnormal or damaged sperm.

4. **Sexual behavior assessment:** Keenness to serve (libido) and ability to serve are important for any breeding male. It is particularly important during this part of the examination to watch carefully for any penile deviations which may prevent intromission (the penis entering the vagina) and for evidence of an ejaculatory thrust.

### **Genetic and Venereal diseases**

Periodic physical examination of any candidate or serving bull / buck is needed to ensure that the bulls do not display clinical symptom(s) of any infection or any contagious disease.

Testing for Tuberculosis and Brucellosis should be mandatory every six months. Ideally, bulls should be karyotyped to rule out chromosomal defects. For exotic and crossbred bulls, specific tests may also be conducted for ruling out genetically transmitted diseases like Bovine Leukocyte Adhesion Deficiency (BLAD), Citrullinemia and Deficiency of Uridine Monophosphate Synthase (DUMPS) and Complex Vertebral Malformation (CVM). Vaccination for known viral/bacterial diseases should be compulsory and vaccination records should also be maintained.

### **Maintenance of breeding records**

Service records will be required to be maintained by the bull service provider in the form of register to be periodically verified by designated veterinarian from the Authority and these records must be communicated electronically with the Authority for maintenance of a database through the breed associations as frequently as desired. Moreover, bulls should not be allowed to remain in an area for more than 3 years and in case of rams and bucks, for 1 year, to avoid inbreeding in the population.

**2.IV : Chapter 4**  
**Standards Operating Procedures**  
**For**  
**Physical Infrastructure for Semen Production Facilities**

Physical infrastructure and general management conditions at the semen production and freezing facility can easily affect the quality of semen and consequently the whole livestock value chain if minimum standards are not ensured. Any semen production facility will be required to be registered with the Authority. These will be required to follow minimum standards laid down here.

**1. Premises**

Premises for semen production facilities should be away from public areas with minimum traffic and flow of people. Sufficient trees shall be planted and lawns prepared around the semen station to reduce dust and heat in the summer.

**2. Housing**

There should be provision of animal housing for individual bulls, a semen collection yard, a semen processing room and a semen storage room. For equipment cleaning/ disinfection / sterilization there should be a separate room. Bull sheds shall have spacious individual pens with adequate loafing area, manger and water trough with access to drinking water all time. Adequate shade around the bull shed shall be provided. The roof shall be made of asbestos or some other suitable material. During summer, cooling system with sprinklers and fans may be required particularly for the buffaloes and exotic bulls.

At one corner of the farm, there shall be an isolation shed for separating ailing / sick bull(s) for treatment. Bull(s) once diagnosed suffering from infectious diseases shall be removed immediately from semen station for safety of other bulls. There should be separate staff and separate bio-security arrangements for semen station and quarantine facilities where fresh entrants will stay for 1-2 months. Entry of unauthorized persons must be prohibited.

**3. Management of Bulls**

A satisfactory state of cleanliness of barns and bulls should be ensured. For proper management of bulls, the following points shall be considered:

- a) The bulls shall be kept under hygienic conditions at all times.
- b) The coat of the bulls shall be kept clean and generally short. The hooves shall be regularly trimmed.
- c) The length of the tuft of hairs at the preputial orifice, which is invariably soiled, shall be cut to about 2cm. The hair would not be removed altogether, because of its protective role. If cut too short, it may cause irritation of the preputial mucosa.
- d) Bulls shall be brushed and groomed regularly, and where necessary, special attention shall be given to the underside of the abdomen, a day prior to semen collection.
- e) Cleaning of the prepuce with sterile normal saline solution may be done every ten days if the microbial load is within the prescribed limits. Cleaning prior to the day of collection can be practiced if the microbial load in frozen semen is beyond the prescribed limit.

- f) In the event of obvious soiling, careful cleaning of the preputial orifice and the adjoining areas with soap or a detergent is recommended; followed by thorough rinsing and drying.
- g) Scientific feeding schedule shall be followed for the bulls. Semen station shall carryout routine quality analysis of feed and fodder for arriving at a balanced ration.
- h) Vaccination schedule should also be strictly followed.

#### **4. Conditions for Semen laboratory**

- a) The premises and inside of the laboratory should always be neat and clean. The ceiling and walls of the laboratory shall be made up of non-porous materials. All cracks and crevices shall be sealed to control pests and insects.
- b) Entry of persons to the laboratory, other than laboratory personnel, shall be strictly restricted. Airlock system or anti-room shall be provided to avoid direct entry to the semen-processing laboratory.
- c) Laboratory windows shall preferably be made of double sheet glass with fixed aluminium frame. The glass panes shall be plastered with sun control films to avoid direct sunlight. The doors shall be kept closed, especially during dilutor preparation and semen processing.
- d) Preferably cassette type or, split type air conditioners fitted with air purifying system with remote temperature control mechanism should be installed to maintain the room temperature at 20°C - 22°C. The number of ACs to be fixed to sustain this temperature shall depend on the size of the processing room. Maintaining this temperature is most important to achieve the best results when single step dilution method is followed for freezing semen. The flow of air from AC must not be towards the front side of the Laminar Air Flow Unit. Adequate number of thermometers shall be kept in a few places in the laboratory to check the room temperature. Alternatively, central cooling with 10 to 15 air exchanges should be fixed, especially for the semen processing laboratory. This helps to control the bacterial load in the semen- processing laboratory and in removing obnoxious odour. The processing laboratory should ideally maintain around 55% relative humidity.
- e) Sink drains shall be decontaminated routinely with a disinfectant. Sink shall not be placed in the semen processing room.
- f) The floors shall be preferably made up of vitrified tiles. Floors and horizontal surfaces shall be cleaned and mopped with a disinfectant solution, as dirt and dust, which settle on these surfaces, are the main sources of contamination.
- g) Unwanted furniture, equipment and materials shall not be kept in the laboratory as they only provide additional area for dust and spores to collect.
- h) Appropriate number of germicidal UV lights with respect to area of laboratory, laminar airflow unit, apron and laboratory footwear cabinet may be fixed with a common operating switch outside the laboratory. These lights shall be switched 'on' at least 8 hours prior to commencement of work in the laboratory and shall be switched 'off' before beginning work. The date of installation of the UV lights shall be noted to facilitate replacement as the life of UV tube is limited. A logbook should be maintained for timely replacement of UV lights.
- i) The laboratory shall be fumigated twice a week with a suitable, using humidifier.

- j) Fumigation should be supported by monitoring laboratory environment by bacterial load test. The bacterial load shall be measured every week to monitor pollution of the laboratory atmosphere.
- k) The work platform, the parts of equipment and other items to be handled during processing of semen, shall be cleaned with suitable disinfectant. It is advisable to repeat cleaning schedule after completing processing of semen.
- l) Clean laboratory footwear, apron, hand gloves, mask and caps shall be compulsory put on while working in the laboratory and extra clothing should be removed before entry.
- m) Eating, drinking, smoking, etc. Shall be prohibited in the laboratory and unnecessary conversation should be discouraged. Besides, entry of persons shall be strictly restricted.
- n) Long exposure of semen to ultraviolet rays, visible light in direct sunlight and white florescent light causes chromosomal damage and hence, direct exposure to such sources of light shall be avoided. Hence, there shall be provision for indirect or diffused lighting inside the semen processing room.

## **5. Personnel Hygiene**

Clothing, skin and hair of laboratory personnel are the sources of contamination. Hence, all should wear laboratory aprons and footwear all the time while they are in the laboratory. Hands shall be washed with soap and water and rinsed with 70% alcohol, before commencing work in the laboratory. The bull attendants must undergo test for TB every year. Other staff working on the station should be tested for TB once in two years. Restricted entry inside the semen processing room and freezing room shall be strictly adhered to.

## **6. Equipment**

- a) The exteriors of all equipment and furniture shall be cleaned weekly. The equipment shall be kept covered by plastic covers when not in use.
- b) The pre-filter of Laminar Airflow unit shall be cleaned weekly. Routine servicing and testing twice a year will ensure efficiency of air filters. Alternatively, culture plate test shall be carried out at frequent interval to assess bacterial load of the air passing through the filters.
- c) Digital photometer/Computer aided Spectrophotometer shall be validated with Haemocytometer readings for sperm concentration twice a year separately for cattle and buffalo (20 samples each).
- d) The automatic semen straw filling and sealing machine shall be thoroughly cleaned, immediately after use.
- e) The microscope lens shall be gently cleaned daily with a piece of cotton soaked in a mixture of ethyl and methyl alcohol (1:1) or a mixture of 80% ethyl alcohol and 20% ether).
- f) The bio-freezer shall be defrosted, thoroughly cleaned and dried, immediately after use.
- g) Incubators to maintain artificial vagina shall be cleaned and disinfected with 70% alcohol.
- h) Single distilled water shall be used in autoclave and thermo-controlled water bath. The water bath shall be cleaned and filled with single distilled water on a regular basis.
- i) The thermometer kept immersed in water bath shall be cleaned daily to have precise temperature reading or water bath fitted with digital display temperature indicator should



be used.

- j) The Liquid Nitrogen containers returned / received from foreign countries and contagious disease prone areas shall be disinfected thoroughly.
- k) The refrigerator meant for storing eggs, antibiotics and buffer shall not be used for storing vaccines and other materials. All such materials shall be stored at a place away from semen laboratory. The refrigerator used for storing eggs, etc. shall be sterilized every week using alcohol swab.
- l) All equipment used in semen processing should be covered under annual maintenance and calibration contracts.

### **List of equipment**

- |                         |                                     |
|-------------------------|-------------------------------------|
| 1. Standard Thermometer | 11. Refrigerator(s)                 |
| 2. Water Bath           | 12. Laminar Air Flow Units          |
| 3. Weighing Balance     | 13. Biological Freezer              |
| 4. Incubator            | 14. Microjet Ink Printer            |
| 5. Autoclave            | 15. Straw Filling & Sealing Machine |
| 6. Microscope           | 16. Water distillation unit         |
| 7. Slide Warmer         | 17. Semen tank (large and small)    |
| 8. Micropipettes        | 18. Computer, etc                   |
| 9. pH Meter             | 19. Hot Air Oven                    |
| 10. Photometer          |                                     |

## **2.V: Chapter 5**

### **Standards and Procedures** **For** **Semen Collection and Preservation Standards**

Semen collection is generally done in the morning hours. Ideally, the floor of the collection yard shall be made of concrete layer at a depth of one foot from the ground level. It should be comforting for proper footing of the bulls. Good quality rubber mat (with interlocking arrangement) or coir mat may be put on the mounting area for adequate cushioning effect. After collection, the area must be thoroughly cleaned and disinfected. A dusty floor shall be avoided to prevent dust falling on the AV/ semen samples. On the day of collection, before collecting semen, the bulls shall be properly washed and cleaned. After that, the prepuce shall be cleaned externally with normal saline and a sterilized paper napkin or sterilized cloth napkin soaked in normal saline to remove any sand or dust particles. For each bull a separate napkin shall be used.

The person responsible to carry out preputial wash must use disposable gloves and separate sterilized nozzle for each bull to avoid transmission of infection from one bull to another. Semen collection should be individualized based on the bull. Sexual preparation (number of false mounts and restraint) of the bulls may be done considering the individual behavior of the bulls and not generalized. For this purpose, the sexual behavior of the individual bulls shall be studied and documented. As a general rule, bulls shall be sexually prepared by giving two / three false mounts followed by restraint. The gap between two ejaculates shall be half an hour to one hour depending on the bull. Second ejaculate shall be taken with proper preparation of bulls.

Sterilized bull aprons shall be used to avoid penis touching hindquarter of the dummy. Before every collection, the semen collector shall either wash his hands with a disinfecting solution or use disposable gloves or do both. The semen collector shall not touch the penis.

Preferably veterinarians shall perform semen collection. If semen is collected by staff, a veterinarian shall remain present to supervise the collection process. While taking collection, it shall be ensured that AV is not thrust on penis of bull, instead penis should be guided to AV. Immediately after collection, and the AVs shall be thoroughly cleaned by non-spermicidal neutral detergent. Separate AVs shall be used for each ejaculation.

The AV shall be changed even if the bull has inserted its penis without successful ejaculation. The same AV shall not be used twice. The AVs shall always be kept inverted and the collection tube shall be covered with felt / water jacket (plastic bottle filled with warm water at 37° C) to avoid cold shock. The open end of sterilized AVs shall be covered with aluminum foil, which would be removed at the time when bull is ready for giving semen.

Appropriate size AVs, ranging from 8-14", shall be used only for Cattle & Buffalo to ensure semen is ejaculated in cone. The cone shall be of top quality Neoprene rubber. Use of lubricant shall be avoided. If it is extremely essential to use lubricant, separate sterilized glass rods shall be used for smearing the Jelly on each AV. The AV shall not be shaken after ejaculation; otherwise lubricant and debris may mix with the semen samples. As soon as the first

ejaculate is taken, the bull apron should be removed and dipped in the plastic tub filled with detergent lotion. For second ejaculate, a fresh apron should be tied to the bull.

The entry of visitors and staff/ labourers (other than those not involved in semen collection) shall be strictly prohibited in the collection arena at the time of semen collection. Protective clothing(barn coat) and gumboots shall be used by the veterinarians and personnel during semen collection. Gumboots and barn coat should be washed immediately after completion of semen collection work. Semen stations may follow the norm of minimum two ejaculates per collection and minimum two collections per bull per week for taking 180 ejaculates annually from each adult bull. However, a maximum number of collections per bull would depend on the individual capacity of the male of other species.

### **Semen Evaluation**

Semen evaluation, freezing and packing are critical quality control segments. As soon as the neat semen is received from the collection area, it shall be kept in the laboratory in a thermo-controlled water bath at 37°C under aseptic conditions (such as Laminar Air Flow Unit) and semen volume should be recorded. After examination of sperm concentration and initial motility, semen samples shall be primarily diluted with dilutor maintained at 37°C. After initial dilution of semen in the ratio of 1:1, the semen should be extended further after 7 minutes of cooling at 20°C with dilutor maintained at the lab temperature. The semen samples should not get accumulated for long time in water bath, which may reduce their viability. Sperm concentration shall be checked preferably by a digital photometer with auto dilutor, manufactured by a reputed company. The photometer shall be calibrated separately taking 20 readings each for cattle and buffalo semen, at least once in six months, with haemocytometer readings. Semen samples showing less than 500 million / ml sperm concentration shall be discarded. The volume of straws should be determined as it may vary from batch to batch. While determining the dilution rate as per the photometric reading, the actual volume of mini straw should be fed to the CASA. Straw volume of randomly drawn straws from a day's production should be checked as part of quality assurance and documented. Appropriate antibiotic must be added to the extender.

Semen samples selected for freezing should have minimum 70% initial progressive motility. Final dilution of semen, keeping a minimum of 20 million spermatozoa per dose, shall be done in appropriate flasks with the dilutor maintained at 37°C. Filling and sealing of semen shall be done under aseptic conditions (such as Laminar Air Flow Unit) using sterile straws, filling nozzles and fresh rubber tubing's. French Mini straws should be used.

Unused straws shall be repacked (air-tight) under Laminar Air Flow Unit before storage. Immediately after use, all the glass ware, rubber ware, plastic tips and other re-usables shall be immersed in neutral detergent solution (to be kept in a plastic tub near the Laminar Air Flow Unit). The freezing should be carried out as per the recommended protocols for freezing cattle and buffalo semen. After freezing gets over, the straws should be collected from the racks using scoop tongs. The operator should wear woolen gloves with leather gloves over it to avoid frost injury.

### **Color specifications for different species and breeds:**

All semen stations shall follow the following color codes for filling of semen in straws:

### Printing of Straws

After filling and sealing, information pertaining to supplier (towards left side), bull name, registration number, date of collection, batch number and breed (towards right side) should be printed on straws. Bull registration numbers will follow international norm ([www.icar.org](http://www.icar.org)) and following abbreviations will be used for various breeds. For example for a Semen Production Unit Karachi (Registered with the Authority as (x), semen packed on 28 January 2015 for a Red Sindh bull having popular name as Jumbola registered as (x) with the Authority (bull code), the straw should read as:

Semen Production Unit Karachi PK(X) RS(XX)(Data of Production ) 28,Jan 2015 Red Sindh

#### Colour codes for filling of semen in straws

Breed	Color
Red Sindh	Red
Holstein cattle	Pink
Holstein Crossbred cattle	Light pink
Jersey cattle	Yellow
Kundhi Buffalo	Purple
Goat	Light green
Sheep	Orange
Nili Ravi	Light Grey
Gir	Blue
Rathi	Pistachio Yellow

#### Breed codes for printing on semen straws

Breed / Species	Abbreviation
Kundhi Buffalo	KUB
Red Sindh	RS
Thari	Th
Kankrej	KN
Kamori	KM
Pateri	PT
Holstein	HO
Cross Holstein	XHO
Jersey	JE
Cross Jersey	XJE
Nili Ravi	NI
Gir	GR
Rathi	RT

Straws manufactured by reputed companies are safer to use for production of quality semen. While buying straws, package volume and microbial load in straws shall be checked randomly from the consignment. In addition, some empty straws should be placed in filling and

sealing machine and the machine should be run to see the sealing quality of the straws. In case of any foul smell, it should be presumed that the straws are manufactured from poor plastic which could be toxic to the spermatozoa and can even result in reduced motility on long storage.

### **Post thaw motility**

After freezing, the semen straws shall be stored in a separate container. Post-thaw motility of semen should be examined at 24 hours (after freezing). Differences in observations shall be updated and recorded for the purpose of accepting a particular batch of semen doses. Whenever there is any doubt, post-thaw motility shall be examined by two experienced persons. Preferably, the person involved in evaluation of neat semen, shall not check the post thaw motility. For a minimum concentration of 20 million per dose, minimum acceptable post thaw motility shall be 40%. Semen doses below 50% progressive motility shall be discarded.

### **Quality Checks for frozen semen**

Quarterly testing of random samples from each batch for bacterial load using standard plate count (the standards for acceptable colony forming units (CFUs) in processed semen is 5000 per ml as per OIE norm. If the bacterial load exceeds the OIE limit, the semen doses are to be discarded.) The frozen semen samples should not have uncountable CFUs as they may have pathogenic organisms. Therefore, semen showing crowded CFUs should be subjected to testing for pathogenic organisms by an outside laboratory. Other tests to check the quality of semen may also be beneficial. Validation of photometer shall be done once in 6 months by checking at least 20 samples each for cattle and buffalo. Neat semen shall be examined at an interval of every six months for morphological abnormalities, particularly for crossbred bulls.

Morphological examination of sperms of young bulls must be carried out (at least six samples at weekly intervals) before introducing them in the herd. Semen should not be used if the sample contains a total abnormality of more than 20% and head and mid- piece abnormality (alone) of 7%. Quality checking of semen straws, drawn randomly from the long storage containers once in three months, should be done as a part of quality assurance.

### **Information System**

In order to facilitate the information system, all the bulls maintained by the semen station must be identified by ear tags/ cold branding. The semen stations shall use suitable software to record data pertaining to various activities and also should have online facility for the same. Barcode system can also be used to facilitate traceability.

Volume of semen, density, motility, sperm concentration, dilution rate, total extended volume, post-thaw motility (24 hrs after freezing), and total number of doses produced, etc. shall be maintained. Pre-freeze and post-thaw motility shall be checked for new and problematic bulls.

Miscellaneous information regarding actual reason(s) for not donating semen, undesired percentage of gross morphological defects, semen pH, presence of dirt, dust, blood, pus, etc. in semen samples shall be noted and recorded. Details of semen supplied to various agencies, including post-thaw motility at the time of dispatch, shall be recorded. Fertility data of bulls, conception rate, records of the progeny associated with any genetic defect, percent male / female born, etc. shall be noted and recorded. Report on microbiological examination of semen samples shall also be maintained along with record of all quality tests for neat and frozen semen

samples shall be maintained.

### **Semen Storage**

To avoid accidental spread of diseases, the semen station shall follow the procedure of preserving semen doses for at least 30 days after production. Frozen semen doses produced at least 30 days prior to the date of dispatch should only be supplied for AI. Semen is preserved in liquid nitrogen at  $-196^{\circ}\text{C}$ . Transferring of semen must be done quickly. After checking post-thaw motility, if found acceptable, frozen semen doses shall be kept in temporary storage for 7 days. After temporary storage, the semen goblets shall be transferred to the bulk storage containers with proper recording of position in the canisters. After each dispatch, records redefining the position of remaining doses shall be updated. Frozen semen should be packed in mini goblet (10-15mm) and stored in medium goblet (35mm) and large goblet (65mm). Liquid Nitrogen shall be replenished at regular intervals depending on the liquid nitrogen evaporation rate as per season of the year. The level of liquid nitrogen in the storage tank should not be allowed to go below 1/3rd the depth of the tank. In shipping and handling, the exposure time of frozen semen to air must not be more than 3 seconds. All transfers should be made in mini goblets within 3 seconds.

Two reference samples of the doses dispatched to be drawn and retained for six months or a screen shot of randomly selected sample should be stored and a soft copy of which should be given to the customer. The goblets containing the semen should be well identified and precaution should be taken to see that each goblet has sufficient space for liquid nitrogen. Mini straws need special care and should not be exposed above liquid nitrogen even for a short time (10 seconds) as they get warm faster and any exposure causes irreversible damage to sperm viability.

## **2.VI: Chapter 6**

### **Guidelines and Standards** **for** **Import of Exotic Semen**

Import of exotic semen indigenous restriction need must conform to breeding policy which requires development of indigenous breeds of various species to meet the indigenous and export demand of various livestock products and services. Progeny tested cattle semen is however, allowed for crossing with non-descript cattle both for dairy and beef purposes. Import of semen for species other than cattle for experiment (or other purposes) will require special permission.

#### **1. Dairy Cattle**

Breeds: Holstein, Jersey, Gir, Rathi and Red Sindhi

Form of semen: Frozen in liquid nitrogen (in plastic straws)

#### **Bull's attributes:**

1. Origin should preferably be (but not restricting to) United State of America, Canada, Germany, Australia and United Kingdom other countries only when average performance of daughters meet the laid down criteria.
2. For Holsteins, average lactation milk yield of daughters in a standard lactation of 305 days should be at least 10500 liters for North American, 8500 liters for European and 6500 liters for Australian setups. Daughters under multiple production systems should have average milk yield equivalent to European production setups. Average lactation fat of 3.7% or better and average protein 3.0% or better.
3. For Jerseys, average lactation milk yield of daughters in a standard lactation of 305 days should be at least 7500 liters with average fat% of 4.8% or better and protein, 3.6% or better. These averages may vary and will be kept close to yearly averages reported by International Committee for Animal Recording ([www.icar.org](http://www.icar.org)) for various countries.
4. Should have at least 50 daughters performance recorded in 20 or more herds with better than 80% accuracy of production traits. If genomic information is added to recorded performance, accuracy will increase.
5. Estimated Breeding values or Predicted Transmitting Abilities (Genomic or otherwise) or for milk, fat and protein yields should be positive in the most recent genetic evaluation. Semen from negatively ranked bulls for production traits is not allowed.
6. Bull should be improver for overall Type (positive PTA for overall Type) alongwith positive PTAs for Udder and Feet & Legs conformation.
7. Bulls should not cause more dystokia than average bulls in the recent most genetic evaluation nor should their daughters prone to mastitis than the average bulls.
8. Bulls should preferably be A2A2 genotype for Beta casein.

9. Bull should be free from genetic disease such as Bovine Leukocyte Adhesion Disease, (BLAD) Deficiency of Uridine Monophosphate Synthetase (DUMPS), Complex Vertebral malformation (CVM), Citrulinemia, Factor XI etc and should have normal karyotype.
10. Should be free from scheduled reproductive problems transmittable venereal diseases duly certified by the concerned department of the country. Health certificate from competent authority of the country of origin will have to be supplied.
11. The motility percentage of sperm in frozen semen should not be less than 50% after thawing of semen.
12. Each semen straw (0.25/0.5 ml capacity) should carry at least 20 million sperms for non-sexed and 2 million for sexed semen. Accuracy of female births should be 90% or better with sexed semen.
13. Should be registered with and traceable from Breed Association / Semen Supplier Website.

## **2. Beef cattle**

Breeds: Angus, Hereford, Charolais, Brahman, Simmental, Nelore (panted), Nelore (White) and Kankrej etc (for export of beef from crossbreds) Form of semen: Frozen in liquid nitrogen in plastic straws

### **Bull's attributes:**

1. Origin should preferably be (but not restricting to) United State of America, Canada, Germany, Australia, Brazil and United Kingdom
2. Must be registered with relevant Breed Association
3. Expected Progeny Difference (or breeding value) should be positive (above average) for weaning weight, yearling weight, calving ease and dressing percentage.
4. The accuracy value for various traits should be greater than or equal to 70%.
5. The sperm concentration per frozen dose of semen should be at least 20 million in 0.5/0.25ml straws The motility of sperms in after thawing of frozen semen should be better than 50%.
6. The donor bull should be certified not to have any genetic and venereal diseases.



## **2. VII: Chapter 7**

### **Standards and Procedures** **For** **Artificial Insemination Training and Training Institutions**

Artificial Insemination Training Institutes operating in Sindh will have to register with the Authority and get accredited for issuing AI training certificates. This also applies to any NGO or Government institution involved in the business. Quality of AI training varies across the organizations due to absence of a uniform training module, standard protocol and a mechanism to ensure its effective implementation by the training institutes. The standards are basic minimums of training for undertaking artificial insemination services successfully.

#### **A. Qualification, duration and Curriculum**

##### **1. Qualification and duration**

- i. Diploma of Stock Assistant
- ii. Six months (out of which at least 3 months should be extensive practical training in the field)

##### **2. Curriculum**

###### **a. Theory**

- 1) Different breeds of cows and buffaloes and their production and reproduction parameters.
- 2) Introduction to AI, and its importance, role of AI in genetic up gradation.
- 3) Natural Service (NS) vs. AI, advantages and limitations.
- 4) External and internal body parts of a dairy animal and their functions.
- 5) Male reproductive organs & their functions.
- 6) Semen, its collection, evaluation, processing, preservation.
- 7) Reading a semen straw.
- 8) Female reproductive organs & their functions.
- 9) Important reproductive hormones and their role in reproductive cycle.
- 10) Normal reproductive cycle.
- 11) Oestrus cycle and signs of estrous.
- 12) Puberty, Maturity, Breeding, Fertilization, Implantation, Gestation and Calving.
- 13) Service period, calving interval, and dry period and gestation period.
- 14) Synchronization of breeding females.
- 15) Process of insemination using liquid and frozen semen.
- 16) Importance of fixed time AI.
- 17) Care of animals during & after insemination.
- 18) AI equipment and accessories & their care.
- 19) Liquid nitrogen handling.
- 20) Significance of early pregnancy diagnosis.
- 21) Ultrasonography.
- 22) Methods of calculating conception rates and factors affecting conception rates.

- 23) Common reproductive disorders/ diseases, repeat breeding, causes of abortion, etc.
- 24) Measures to obtain maximum fertility.
- 25) Ear tagging, importance of record keeping, recording formats and submission of records using data transfer devices and web services.
- 26) Method of non-surgical castration.
- 27) Care and management of new born calf and heifers till it becomes pregnant.
- 28) Care and management of animals before and after calving, precautions at the time of calving and use of naval kit for disinfection of naval cord.
- 29) Importance of bio-security measures to be adopted during A.I.
- 30) Basic aspects of nutrition.
- 31) Importance of proper nutrition including feeding of vitamins, mineral mixtures and deworming in fertility management.
- 32) Body condition (scoring) and its relationship with fertility.
- 33) Conservation and development of indigenous breeds through selective breeding.
- 34) Benefits of crossbreeding and genetic improvement of dairy animals.
- 35) The existing Breeding Policy and its enforcement.
- 36) Various government schemes in the livestock sector.

#### **b. Practical**

- 1) Identification of different female reproductive organs on morbid genitalia.
- 2) Palpation of female genitalia in a Phantom box and passing of AI gun.
- 3) Structure of LN container:
  - Different models.
  - Handling & care.
  - Filling and checking LN level.
- 4) AI equipment & accessories:
  - Handling & care including sterilization.
  - Reading and recording of semen straw information.
- 5) Handling of animals under different situations.
- 6) Palpation of female genitalia in live animal.
- 7) Passing of AI gun in live animals.
- 8) Reproductive examination of cows/buffaloes in estrus.
- 9) Demonstration and extensive practice of:
  - Proper method for withdrawal of straw from containers.
  - Proper thawing procedure.
  - Proper preparation of gun.
  - Correct site of semen deposition.
- 10) Demonstration of semen motility as affected by long time exposure and thawing process.
- 11) Introduction of metritis and its basic treatment.
- 12) Pregnancy diagnosis at 40 days & beyond.
- 13) Assessing the stage of fetal life.
- 14) Ear tagging.
- 15) Record keeping and communication
- 16) Study visits should be arranged to AI Centre and Semen Production and Processing facility with a visit of any modern Dairy Farm / Feed factory/ fodder farm.

### **3. Tests and examination:**

- Two quarterly written tests on topics covered (10% each)
- Final written test at the end of 6-months (30%)
- Final practical test to evaluate the skills learnt (50%)

### **4. Pass marks:**

- Minimum 50% each in theory and practical

### **5. Attending of in-service short training course of 1-2 days (every year)**

## **B. Physical facilities**

### **1. Class room facilities:**

For a batch of 25 trainees, a class room having minimum of 400 square feet area will be required. If there are more than 25 trainees, there should be an additional class of 400 square feet area. A laboratory having minimum 500 square feet area for practical classes will also be required. This laboratory should have facility to store reproductive organs and keep different models of animals and reproductive organs and space to keep semen and liquid nitrogen storage containers. Disposal of wastes should also be appropriate to keep the environment neat and tidy. There should be a library/reading room having books and journals on cattle, breeding, indigenous breeds and dairy apart from rooms for teachers and administrators.

### **2. Teaching aids**

#### **a. The class rooms must have the following:**

- Adequate chairs and tables for trainees.
- White board.
- LCD Projector.
- Computer.
- Charts and Models.
- The centre must have the required quantity of semen doses and LN storage containers, AI guns and other AI accessories.
- Reproductive organs must be obtained from a nearby slaughter house for palpation and passing AI guns.
- Ear tags and ear tag applicators.
- Measuring tapes and rulers for estimation of heights and body weight.
- Other communication technology aids (notebooks, android phone, printers etc).

#### **b. Animals and housing facilities for practical training**

- For practical's, the centre should have at least one animal for every 3-4 trainees. Alternatively, centre may have contractual arrangement with some culled animal facility, slaughter house or any other farm for practical training. Every trainee must pass AI gun in at least 20 live animals during entire period of class room training before going for field training during later half of the course. Animals used for training should have normal genital tract. Pregnant animals at various stages pregnancy should also be arranged for demonstration.
- If the centre has its own animals, there should be kept in a comfortable shed and fed, watered and vaccinated properly. Restraining facilities for individual animals should also be appropriate. These experimental animals should be replaced every six months. Disposable gloves, dungarees

and gumboots are expected to be worn by working with animals including the trainees.

### **3. Lodging and boarding facilities for trainees**

- The centre should have proper residential facilities for trainees including kitchen and minimum recreational facilities. However, if this not feasible, boarding and lodging facilities from other agencies can also be hired. Copy of the formal agreements with such agencies should be available for record for requirement at the time of Accreditation process.

### **4. Field training**

- The Centre should have some formal arrangement with AI service providing organizations for its trainees to receive apprenticeship training for at least 90 days. During this period, each trainee should do minimum 75-100 AIs and the same numbers of pregnancy tests.

### **5. Documentation of AI training activities**

- Trainees' records of registration, daily attendance, list of passing out trainees, feedback files, training materials and annual progress reports should be available with the centre.

## **C. Faculty profile and requirement (for a batch size of 25 trainees)**

### **1. Veterinary Officers:**

Minimum two Veterinarians/Teachers registered with PVMC having 3 years of work experience in AI, breeding, health and management of dairy animals along with experience in providing on the job practical training and delivery of lectures.

### **2. Support Staff:**

Minimum one support staff is required with graduation in any discipline

### **3. Labour for cleaning and other services.**

Number should be sufficient to keep the facilities neat and clean and well guarded as well.

## **D. Accreditation of Training Institutes**

### **Each accredited AI Training Institution needs to:**

Follow the standard curriculum and duration for class room and practical training as described in above.

- a) Have arrangements for class rooms with teaching aids and hands on training in palpating morbid reproductive organs in the laboratory and in live animals and for carrying out artificial insemination as mentioned above
- b) Have arrangements with AI Service providing organizations to provide practical training to its trainees for the duration approved
- c) Have a registered veterinary practitioner under whose supervision the institute imparts training.

d) Evaluate the trainees and award a certificate after successful completion of the course. Furthermore accredited institutes will send the names of the successful trainees to the Authority for issue of a license bearing a photograph and a registration number of the trainee. On successful training the training institute shall equip each trainee the AI kit as well. The process of accreditation will start by a written request of any institution intending to award AI training. Submission of application should accompany the required documents which will be verified by the Authority against prescribed Minimum Standards detailed above. Following documents are needed:

- Mission & Objective of AI training Institute.
- Memorandum of Article/ Bylaws of the Organization/AI training Institute as applicable.
- Details of Infrastructure (Office, Classroom, Board & Lodging, etc.)
- Learning resources /teaching aids.
- Facilities available for practical and on the job field training on animals.
- Details of training programmes, duration, and course curriculum, training schedule, admission norms and internal evaluation process.
- Details of training faculty.
- Copies of records maintained by the AITI regarding previously trained persons (names, addresses and contacts).
- Progress for the last two years (except for the 1 time).

In case application is accepted, the application is further processed for a visit by a team and upon verification of required minimums, Accreditation certificate will be awarded to the applicant for a period of two years. For renewal of accreditation, the AITI has to reapply and same procedure will be followed. The AITI must respond to any queries of the Authority as and when asked. Once, accredited, annual progress reports have to be submitted before the end of July every year.

All the AI technicians working in the field before these laws have to go through a refresher course (to be arranged by accreted institutions) to get practicing license from the Authority.

## 2. VIII: Chapter 8

### **Guidelines and Standards** **For** **Use of Exotic Semen**

Exotic semen of cattle is imported to upgrade non-descript local cattle for dairy and beef purposes. Semen of following breeds is allowed for such crossbreeding projects/schemes.

#### **Dairy cattle:**

1. Holstein Friesian
2. Jersey

#### **Beef cattle:**

1. Angus
2. Hereford
3. Charolais
4. Brahman
5. Simmental

For dairy cattle, restricting semen to few dairy cattle breeds is basically to manage the availability of semen from crossbred bulls of these breeds. This is in conformity to the breeding policy where two internationally recommended breeds Holstein and Jersey have been allowed. Even with just two breeds, four types of semen will be needed to keep the desired level of exotic inheritance between 50 and 75% i.e two purebreds and two 50% crossbred bulls. Semen from Brown Swiss has also been imported previously but experience from other countries with similar production system as that of Sindh, Holstein and Jersey should be sufficient. It may also be mentioned that Friesian being a large sized breed is suggested for crossing non-descript when cow being inseminated has a larger size and production system is not subsistence low input type Jersey semen should be preferred for breeding smaller sized cattle. Yet, in both cases production system should be at least medium input system.

For beef cattle, 4 allowed breeds (Angus, Hereford and Charolais, Simmental) have previously been tested for producing crossbred beef animals in Sindh. Addition of Brahman is for production of beef from crossbred cattle for export purpose (beef qualities almost similar to our local cattle). Cows selected for crossing with the exotic breeds should be of bigger size because genetic potential for birth weight of all these cattle is almost double than our local non-descript cattle. The precise breeding guidelines will be as follows:

#### **Dairy cattle**

1. Purebred Red Sindhi and Red Sindhi like cows – should be bred only to Red Sindh semen.
2. Pure breed Thari and Thari like cows should be bred only to Thari.
3. Pure Kankrej and Kamkraj like cows should be bred only to Kankrej.

4. Crossbred cattle having 75% or higher exotic inheritance should be bred with 50% exotic bulls.
5. Crossbred cattle having less than equal to 50% exotic inheritance – may be bred with purebred exotic semen for high input systems and with 50% crossbred bulls for low to medium input systems.
6. Non-descript cattle (smaller size) – should be bred with Jersey semen but under low-medium input system, Red Sindhi or Thari may be used as well.
7. Non-descript cattle (bigger size) may be bred with Holstein or Jersey semen (depending upon the input system, high input production system may prefer Holsteins).

#### **Crossbreeding local cows with exotic bull/semen for dairy purpose**

	<b>Bull/Semen</b>						
<b>Cow</b>	<b>RS</b>	<b>TH</b>	<b>KN</b>	<b>HO100</b>	<b>HO50</b>	<b>JE100</b>	<b>JE50</b>
Red Sindhi (RS)							
Thari (TH)							
Kankrej (KN)							
Crossbred $\geq 75\%$ exotic							
Crossbred 50-75% exotic							
Crossbred $< 50\%$ exotic							
Non-Descript1 (ND1)							
Non-Descript2 (ND2)							

ND1 means smaller sized non-descript cattle, upgraded with Red Sindhi or Thari.

ND2 means bigger sized non-descript cattle.

HO means exotic Holstein (50, 100 is proportion of exotic level).

JE means exotic Jersey.

Males produced from all these crosses should be culled and only top class males be retained (by planned mating) from top class cows for production of future sires of various exotic level.

#### **Beef cattle**

Crossbreeding for beef cattle is not generally recommended for day to day beef production. This is mainly due to high cost of inputs especially when millions of surplus males are available in the province for fattening. However, this may be economical for export (both veal and beef). Another use may be producing of sacrificial animals for big city markets. Following mating plan is suggested assuming that crossbreds (both males and females) will be consumed at a suitable age for sacrificial purpose or for export.

**Crossbreeding local cows with exotic bull/semen for beef purpose**

<b>Exotic semen</b>	<b>Scope</b>	<b>Local cows</b>
Angus	Good scope for sacrificial local markets	Nondescript and low yielding crossbred cows
Hereford	Good scope for sacrificial local markets	Nondescript and low yielding crossbred cows
Charolaise	For leaner beef export markets	Nondescript and low yielding crossbred cows
Brahman	For gulf markets requiring Zebu beef	Nondescript local cows
Simmental	For leaner beef Export	Non-descriptive and low yielding



## **2. IX: Chapter 9**

### **Standard Operating Procedure** **for** **Artificial Insemination Services**

Genetic potential of AI bulls is expected to be better than those used for natural service. Risk of transmission of venereal and genetic diseases is also lower yet, conception rate is generally low which can be greatly affected by AI technicians and AI service providers.

The first and foremost issue is clarity regarding the breeding policy of the province. While crossbreeding of cattle is allowed but for non-descript and crossbreds cattle only. Only two breeds are allowed, Holstein and Jersey and desired level of exotic inheritance should be within 50 and 75% range. Holstein Friesian is not preferred for short sized cattle not it is for low input system / poor farmers. Purebreds cattle such as Red Sindhi and Thari (and cows which as almost similar to them) should be bred with Red Sindhi and Thari cattle semen only. Kundhi buffaloes should be bred with Kundhi buffalo semen. Similarly, Tharparkar, Red Sindhi, Kankrej strains should be bred with respective bulls or semen from them. Same shall be true for sheep. The general slogan is: purebreds for purebreds and exotics and crossbreds for non-descript and crossbreds.

The second issue is availability of semen for Authority registered semen production facilities only. Top graded semen production facilities should be preferred.

#### **Semen storage**

- Store frozen semen doses in a well-ventilated, all weather safe storage area.
- Ensure a proper and foolproof identification system for each semen container, canister, and goblet so that a bull's semen can be traced with ease.
- While transferring semen doses, goblets should be well identified and precaution should be taken to see that each goblet has sufficient space for liquid nitrogen. Frozen semen should not be exposed above liquid nitrogen as it may cause irreversible damage to sperm viability.
- All transfers of semen straws into goblets should take place under liquid nitrogen, in a polystyrene / thermopol box.
- Liquid Nitrogen should be replenished in both storage and distribution containers at regular intervals to ensure proper level of liquid nitrogen.
- Details of semen doses supplied to various AI technicians at the time of dispatch should be recorded. After each dispatch, records redefining the position of remaining doses should be updated.

#### **Liquid Nitrogen**

Service provider shall have a bulk Liquid Nitrogen (LN) sourcing, storage and delivery facility. A schedule of LN replenishment to all AI centres on fortnightly / monthly /quarterly basis, whichever is convenient depending on the field containers shall be worked out by each service provider and shall be adhered to in the interest of maintaining quality of semen. A log book shall be maintained for all such schedules at different locations/ starting points of supply routes. The AI centre should have a bigger container for LN and semen straw storage, preferably of 35 litre capacity, and a small portable LN container, preferably 2-3 litre capacity, to carry the

semen to the place where the AI is carried out. The LN containers in the AI centre shall be protected sufficiently to avoid damage to container. AI centre should have a dip stick with critical level marks and a ready-reckoner for assessing the LN levels and quantity of LN in litres. Supply of LN should be either through portable LN tankers of 500 to 2000 litre capacities with gravitational flow or through the LN delivery pump and not by pouring LN from one container to another container.

#### **A.I guns and sheaths**

Stainless steel AI guns from an agency whose AI guns are tested and approved by Authority shall be used. Same shall be true for AI. AI accessories like forceps and scissors shall be made of good quality stainless steel. Thermos flask and thermometer shall be of good quality.

#### **A.I Technicians**

AI Service providers should ensure that they engage only those AI technicians who have undergone a training course in AI from a government recognized AI training institute and collect a copy of their training certificate at the time of appointment.

#### **Animal Identification**

Every animal receiving AI shall be identified with an Ear Tag with a unique number and (preferably) a barcode. These numbers shall enable generation of reports concerned to the individual animal and the associated information through an information system. Only polyurethane laser printed ear tags having a 12 digit number and a bar code shall be used. The numbering system followed shall be unique with the last digit of the number being a “check digit” to ensure that no two animals are tagged with the same number. Only numbers supplied by an agency identified by the Authority shall be used for unique identification of animals. An example may be for a buffalo tagged in Karachi district (digit code 16) and registered as 000001 may have ear tag as 16KUB000001.

#### **Supervision, Review of Activities and Communication**

There shall be a hierarchy of supervisory mechanism. Every 20 AI technicians should have a Veterinary Officer. Every 10-15 AI centres should have Deputy Director to provide advisory /Advice services. A team of 100 AI technicians, 10 supervisors and 3 veterinarians shall form a District controlled by a Deputy Director/Additional Director. An effective communication network shall be in place for communication among the team members in a given area.

AI technicians, supervisors and veterinary officers of an area shall meet once in a month for a review of technical programme, business transactions as well as or scheduling the extension programmes. There shall be a monthly review meeting at District level involving Deputy Director, veterinary officers and AI supervisors/technicians. Effective supervision is reflected in the accuracy of reporting, fixing the problems faced efficiently and effectively, acceptance of progress records by the system, promptness in business transactions and minimum backlogs.

## CLINICAL EXAMINATION OF BULL

Unit:		
NAME / IDENTIFICATION NUMBER OF BULL		
BREED:		
DATE OF BIRTH:		
REGISTRATION NUMBER:		
1	CLINICAL EXAMINATION (On First day of Quarantine) (Generation health, testis, penis, accessory glands and presence of heredity defects)	
	Date of examination _____ / _____ / _____ Yr                 mo                 day	
	Find:	
	Remarks	
2	ROUTIEN TEST (Do within a month of semen collection)	
	TUBERCULOSIS:	
	Date of Intradermal Injection: _____ date of reading _____	
	<u>BRUCELLOSIS:</u>	
	CFT or ELISA	
	<u>TRICHOMONIASIS: (Three sheath washes at one week interval )</u>	
	Date of first washing: _____	
	Date of second washing: _____	
Date of third washing: _____		
CAMPYLOBACTERIOSIS: (Three sheath washes at one week interval)		
Date of first washing: _____		
Date of second washing: _____		
Date of third washing: _____		
LEPTOSPIROSIS:		
Agglutination test		

3	CERTIFICATION
	<p>I hereby certify that above information is, to the best of my knowledge, true and correct. A clinical examination was performed by me on _____ and _____ and the above described bull was found healthy and free from any infectious diseases.</p> <p>Signature _____ Date _____</p> <p>Name in Capital letters _____ PVMC registration Number _____</p>
4	ENDORSEMENT ( Bull Approval Committee)
	<p>We hereby endorse the certificate as done above and recommend/do not recommend the use of above described full for semen collection for Artificial Insemination purposes.</p> <p>Signature _____ Date _____</p> <p>Name in Capital letters _____</p> <div style="border: 1px solid black; width: 200px; height: 60px; margin-left: auto; margin-top: 20px; display: flex; align-items: center; justify-content: center;"> Official </div>

**BREEDING SOUNDNESS OF A BULL**

Date of Examination .....

Owner:----- Address:-----

(A). **Identification** Name – (Ear tattoo, Ear, tag, Brand No.)

-----

Breed----- Age-----

(B). **Record**

i) Pedigree covering information up to grand maternal and paternal sides

-----

-----

ii) Progeny: No. of herds/No. of daughter - PD- Progeny Testing based on BLUP -----

-----

iii) Proven /Test:-----

iv) A.I / Natural:-----

(C). **General Clinical Examination**

i) Weight ..... Condition Score:

-----

ii) General Health: .....

iii) Teeth and Jaws: .....Eyes: .....Thorx:

.....Abdomen.....

iv) Feet:.....Legs:.....Joints:.....Gait:.....

(D). **Physical Examination of the Reproductive Organs**i) Scrotum:.....Testis<sup>15</sup>:..... Scrotal Circumference..... (cm)

ii) Normalcy

Consistency.....

iii) Epididymides.....Spermatic Chord:

-----

iv) Accessory organs:

Seminal vesicles.....

Ampulla<sup>16</sup>.....Prostate:

..... Bulbourethral:.....

v) Penis and Prepuce:

-----

(E). **Semen Examination**

i) Collection Method:.....

-----

ii) Volume (ml):..... Density (0-6).....

Wave Motion (0-6)..... Motility (%).....

iii) Examination of Serving Behavior:

Serving Ability:

Libido.....Erection (Stiffness.....Protrusion.....Deviation.....)

Seeking.....Ejaculatory Thrust.....Protrusion of thrusting.....

Body Position.....Serving Capacity.....acceptance of Vagine.....

Mating & semen ejaculation:.....

(F). **Vaccination Record**

Foot & Mouth:.....

Black Quarter:.....

HS:.....

(G). **Tests for infectious Diseases:-**

Brucellosis.....

Leptospirosis:.....

Trichomoniasis: .....

Vibriosis: .....

IBR: .....

BVD” .....

(H). **Semen Result Sheet**

Concentration (million per ml):.....

% spermatozoa alive (Nigrosin eosin):.....

% abnormal sperm heads:.....

% Spermatozoa with:

Proximal cytoplasmic droplets .....

Distal cytoplasmic droplets .....

Tailless heads .....

Coiled tails .....

Acrosome defects .....

Structural abnormalities .....

Cells other than spermatozoa .....

**Comments:**

**BRUCELLOSIS TESTING**

I	Screening Test Details	Name	ELISA, RBPT, CFT, SAT
		Sample	Serum
		Testing at	CVDL, or any other
II	Eligible animals		<ul style="list-style-type: none"> <li>• All above one year</li> <li>• In females 14 days after calving or abortion</li> </ul>
III	Frequency of Testing	Positive herd	30 to 60 days after culling of last positive animal
		Negative herd	Exactly one year ( $\pm$ 1 week) after last whole herd negative testing
		Negative herd (Option	Where the diseases have been maintaining a very low profit (less than 1% positive) quarterly or six monthly sample could be collected to minimize losses.
IV	Action on finding a positive bull	Animal	Immediate isolation, castration and culling.
		Semen	Destroy semen doses since last negative test.
V	Brucellosis Free herd (OIE)		Herd found negative on two consecutive annual tests
VI	Quarantine	Duration of quarantine	Minimum 30 days
		Test schedule	Two test, Serum ELISA, interval of 30 days between tests, only negative animals to be allowed to mix with the rest of the herd.
VII	Additional testing at Sexual maturity		Serum ELISA before bulls are used for semen collection and distribution for AI

CVDL- Tando Jam  
Sindh Agriculture University Tando Jam  
SBBUVAS Sakrand.

**CAMPYLOBACTERIOSIS (VIBRIOSIS) TESTING**

I	Screening Test Details	Name	Bacterial isolation & identification
		Sample	Perpetual washing, Semen
		Testing at	CVDL, Sindh Agriculture University Tando Jam, SBBUVAS Sakrand.
II	Eligible Animal		All male animals
III	Prevention		Annual sheath Washing
IV	Frequency of testing	Positive herd	30 days after culling of positive animal
		Negative herd	Exactly one year ( $\pm$ 1 week) after last whole herd negative testing
V	Action on finding a positive bull	Animals	Treat the animal
		Semen	Destroy semen doses since last negative test.
VI	Quarantine	Test Schedule	One test if age is less than 6 months, others 3 consecutive tests at weekly interval.



**TRICHOMONIASIS TESTING**

I	Screening Test Details	Name	Agent isolation & identification
		Sample	Perpetual washing
		Testing at	CVDL, Sindh Agriculture University Tando Jam, SBBUVAS Sakrand.
II	Eligible Animal		All male animals
III	Prevention		Annual sheath Washing
IV	Frequency of testing		Annual
V	Action on Finding a Positive animal	Animal	Treat animals.
		Semen	Destroy the collections since last negative test.
VI	Quarantine	Duration of Quarantine	Minimum 30 days.
		Test Schedule	One test if age is less than 6 months, other 3 consecutive tests at weekly intervals.
VII	Addition Testing at Sexual Maturity		Protozoa isolation before bulls are used for semen distribution for A.I.

## LEPTOSPIROSIS TESTING

I	Screening Test Details	Name	Agglutination test, immuno-fluorescence, immuno-histochemistry, PCR
		Sample	
		Testing at	CVDL, Sindh Agriculture University Tando Jam, SBBUVAS Sakrand.
II	Eligible Animal		All male animals
III	Prevention		Screening and isolation of animals. Prevention from rodents
IV	Frequency of testing		Annual testing
V	Action on Finding a Positive animal	Animal	Treat animals.
		Semen	Destroy the collections since last negative test.
VI	Quarantine	Duration of Quarantine	Minimum 30 days.
		Test Schedule	Annually
VII	Addition Testing at Sexual Maturity		ELISA, annually

## TUBERCULOSIS TESTING

I	Screening Test Details	Name	Delayed Hypersensitivity Single Intradermal test
		Reagent	Bovine tuberculin PPD
		Testing at	CVDL, Sindh Agriculture University Tando Jam, SBBUVAS Sakrand.
		Positive result criteria	As per OIE norms/ <i>Negative:</i> increase in skin thickens less than 2mm & without clinical signs, viz; exudation, necrosis, pain, inflammation of the lymphatic duct of that region or the lymph node, 72 hours post inoculation.  <i>Inconclusive:</i> Increase in skin thickness more than 2mm & less than 4mm, absence of above clinical signs, 72 hours post-inoculation.  <i>Positive:</i> increase in skin thickens 4mm or more, or presence of clinical signs viz; exudation, necrosis, pain, and inflammation of the lymphatic duct of that region or the lymph node, 72 hours post-inoculation.
II	Eligible Animal		All animals above 6 week of age.
III	Frequency of testing	Positive herd	Minimum 60 days after culling of test positive animals
IV	Action on finding a positive bull	Animal	Immediate isolation and culling
		Semen	Destroy semen doses since last negative test
V	Tuberculosis free herd (OIE)		Herd found negative on two consecutive tuberculin test at an interval of 6 months, the first being performed 6 months after the slaughter of last affected animal.
VI	Quarantine	Duration of Quarantine	Minimum 90 days
		Test Schedule	Two tuberculin tests, minimum interval of 60 days between tests.

PPD-Purified Protein Derivative

## FOOT AND MOUTH DISEASES TESTING

I	Action during FMD outbreak	Animal	Isolate diseased animals till recovery, do not cull.
		Semen	<p>Semen from FMD infected bulls: Destroy semen collected during one month before onset of break. Do not collect semen from bulls during the outbreak and three months after the last case of FMD recovered in the farm infected animals must be given 90 days rest.</p> <p>Semen from healthy bulls maintained in FMD infected farm: Destroy semen collected during one month before onset of outbreak. Do not collect semen from bulls during the outbreak and one month after the last case of FMD recovered in the farm.</p> <p>Semen could be used other than the above mentioned periods, if there is no new case of FMD develops during three months period after last FMD case recovered in the farm.</p>
II	Quarantine		Vaccination – as per manufacture’s recommendations.
III	Vaccination	Oil Vaccine	<p>Annual in farm and 5 to 10 km around the farm</p> <p>Test for seroconversion, by collecting serum on the day of vaccination and 21 days later.</p>

RERGISTRAR

No.

DATED

- - 2017.

A copy is forwarded to the Superintendent, Government Printing Press Sindh, Hyderabad for publication in the Extra-ordinary issue of the Sindh Gazette and supply of 30 printed copies of the notification to this office.

RERGISTRAR

**NO.&DATE EVEN**

A Copy is forwarded for information and necessary action to:-

1. The Chief Secretary, Sindh.
2. The Principal Secretary to Governor, Sindh.
3. The Secretary to Chief Minister, Sindh.
4. The Chairman, Planning & Development Board, Sindh.
5. The Senior Member, Board of Revenue, Sindh.
6. The Secretary, Provincial Assembly, Sindh.
7. Accountant General, Sindh.
8. Vice Chancellors SAU Tando Jam/SBBUVAS, Sakrand.
9. The Director General Extension & Research.
10. Executive Directors Livestock Department.
11. President Livestock Breeder Association
12. Pakistan Dairy Association

**RERISTRAR**